

קורס גרפיקה ממוחשבת

2009 סמסטר ב'

Lighting



מבוסס (מאוד) על


Thomas Funkhouser

Princeton University

COS 426, Fall 2000

Ray Casting


```
Image RayCast(Camera camera, Scene scene, int width, int height)
{
    Image image = new Image(width, height);
    for (int i = 0; i < width; i++) {
        for (int j = 0; j < height; j++) {
            Ray ray = ConstructRayThroughPixel(camera, i, j);
            Intersection hit = FindIntersection(ray, scene);
            image[i][j] = GetColor(scene, ray, hit);
        }
    }
    return image;
}
```



Wireframe

Ray Casting


```
Image RayCast(Camera camera, Scene scene, int width, int height)
{
    Image image = new Image(width, height);
    for (int i = 0; i < width; i++) {
        for (int j = 0; j < height; j++) {
            Ray ray = ConstructRayThroughPixel(camera, i, j);
            Intersection hit = FindIntersection(ray, scene);
            image[i][j] = GetColor(scene, ray, hit);
        }
    }
    return image;
}
```



Without Illumination

Ray Casting

```
Image RayCast(Camera camera, Scene scene, int width, int height)
{
    Image image = new Image(width, height);
    for (int i = 0; i < width; i++) {
        for (int j = 0; j < height; j++) {
            Ray ray = ConstructRayThroughPixel(camera, i, j);
            Intersection hit = FindIntersection(ray, scene);
            image[i][j] = GetColor(scene, ray, hit);
        }
    }
    return image;
}
```




With Illumination

Illumination


- How do we compute radiance for a sample ray?

`image[i][j] = GetColor(scene, ray, hit);`




Goal

- Must derive computer models for ...
 - Emission at light sources
 - Scattering at surfaces
 - Reception at the camera
- Desirable features ...
 - Concise
 - Efficient to compute
 - “Accurate”




Overview

- **Direct Illumination**
 - Emission at light sources
 - Scattering at surfaces
- **Global illumination**
 - Shadows
 - Refractions
 - Inter-object reflections




Shadows



Direct Illumination


Modeling Light Sources

- $I_L(x,y,z,\theta,\phi,\lambda)$...
 - describes the intensity of energy,
 - leaving a light source, ...
 - arriving at location (x,y,z) , ...
 - from direction (θ,ϕ) , ...
 - with wavelength λ






Empirical Models

- Ideally measure irradiant energy for “all” situations
 - Too much storage
 - Difficult in practice




OpenGL Light Source Models

- Simple mathematical models:
 - Point light
 - Directional light
 - Spot light



Point Light Source


- Models omni-directional point source (e.g., bulb)
 - intensity (I_0),
 - position (p_x, p_y, p_z),
 - factors (k_c, k_l, k_q) for attenuation with distance (d)



$$I_L = \frac{I_0}{k_c + k_l d + k_q d^2}$$

Directional Light Source

- Models point light source at infinity (e.g., sun)
 - intensity (I_0),
 - direction (dx, dy, dz)




No attenuation
with distance

$$I_L = I_0$$

Spot Light Source


- Models point light source with direction (e.g., Luxo)
 - intensity (I_0),
 - position (p_x, p_y, p_z),
 - direction $D=(dx, dy, dz)$
 - attenuation



$$I_L = \frac{I_0 (D \cdot L)}{k_c + k_1 d + k_q d^2}$$

Overview


- **Direct Illumination**
 - Emission at light sources
 - Scattering at surfaces
- **Global illumination**
 - Shadows
 - Refractions
 - Inter-object reflections



Direct Illumination


Modeling Surface Reflectance

- $R_s(\theta, \phi, \gamma, \psi, \lambda)$...
 - describes the amount of incident energy,
 - arriving from direction (θ, ϕ) , ...
 - leaving in direction (γ, ψ) , ...
 - with wavelength λ



Empirical Models


- Ideally measure radiant energy for “all” combinations of incident angles
 - Too much storage
 - Difficult in practice



Empirical Models

- Example: BRDF (Bidirectional reflectance distribution function)
 - 4-dimensional function which defines light reflection at an opaque surface.


$$f_r(w_i, w_o) = \frac{dL_r(w_o)}{dE_i(w_i)} = \frac{dL_r(w_o)}{L_i(w_i) \cos(\theta_i) dw_i}$$



OpenGL Reflectance Model


- Simple analytic model:
 - diffuse reflection +
 - specular reflection +
 - emission +
 - “ambient”

Based on model
proposed by Phong in
his PhD dissertation
1973



Diffuse Reflection


- Diffuse: Spread Out / To pass by spreading every way / To extend in all directions
- Assume surface reflects equally in all directions
 - Examples: chalk, clay



Diffuse Reflection

- How much light is reflected?
 - Depends on angle of incident light

$$dL = dA \cos \Theta$$




Diffuse Reflection

- Lambertian model
 - cosine law (dot product)

$$N \cdot L = |N| |L| \cos \Theta$$


$$\hat{N} \cdot \hat{L} = \cos \Theta$$

$$I_D = K_D (\hat{N} \cdot \hat{L}) I_L$$





OpenGL Reflectance Model

- Simple analytic model:
 - diffuse reflection +
 - specular reflection +
 - emission +
 - “ambient”




Specular Reflection

- Reflection is strongest near mirror angle
 - Examples: mirrors, metals



Specular Reflection

- How much light is seen?
 - Depends on angle of incident light and angle to viewer




Specular Reflection

- Phong Model

- $\cos(\alpha)^n$


Phong exponent: apparent smoothness of the surface



$$I_S = K_S (V \cdot R)^n I_L$$

Specular Reflection


- Phong Examples



Direction of light source and shininess exponent is varied

OpenGL Reflectance Model

- Simple analytic model:
 - diffuse reflection +
 - specular reflection +
 - **emission** +
 - “ambient”



Emission


- Represents light emanating directly from polygon

Emission $\neq 0$



OpenGL Reflectance Model

- Simple analytic model:
 - diffuse reflection +
 - specular reflection +
 - emission +
 - “ambient”



Ambient Term


- Represents reflection of all indirect illumination



This is a total hack (avoids complexity of global illumination)!


OpenGL Reflectance Model

- Simple analytic model:
 - diffuse reflection +
 - specular reflection +
 - emission +
 - “ambient”




OpenGL Reflectance Model

- Simple analytic model:
 - diffuse reflection +
 - specular reflection +
 - emission +
 - “ambient”



Surface Illumination Calculation


- Single light source:



$$I = I_E + K_A I_{AL} + K_D (N \cdot L) I_L + K_S (V \cdot R)^n I_L$$

Surface Illumination Calculation

- Multiple light sources:



$$I = I_E + K_A I_{AL} + \sum_i (K_D (N \cdot L_i) I_i + K_S (V \cdot R_i)^n I_i)$$

Overview

- Direct Illumination
 - Emission at light sources
 - Scattering at surfaces
- Global illumination
 - Shadows
 - Transmissions
 - Inter-object reflections



Global Illumination


Global Illumination



"Balanza" © [Jaime Vives Piqueres](#) (2002)

Shadows

- Shadow terms tell which light sources are blocked
 - Cast ray towards each light source L_i
 - $S_i = 0$ if ray is blocked, $S_i = 1$ otherwise




Shadow
Term



$$I = I_E + K_A I_A + \sum_L (K_D (N \cdot L) + K_S (V \cdot R)^n) S_L I_L$$

Ray Casting


- Trace primary rays from camera
 - Direct illumination from unblocked lights only



$$I = I_E + K_A I_A + \sum_L (K_D (N \cdot L) + K_S (V \cdot R)^n) S_L I_L$$

Recursive Ray Tracing


- Also trace secondary rays from hit surfaces
 - Global illumination from mirror reflection and transparency



$$I = I_E + K_A I_A + \sum_L (K_D (N \cdot L) + K_S (V \cdot R)^n) S_L I_L + K_S I_R + K_T I_T$$

Mirror reflections


- Trace secondary ray in direction of mirror reflection
 - Evaluate radiance along secondary ray and include it into illumination model



$$I = I_E + K_A I_A + \sum_L (K_D (N \cdot L) + K_S (V \cdot R)^n) S_L I_L + K_S I_R + K_T I_T$$

Transparency

- Trace secondary ray in direction of refraction
 - Evaluate radiance along secondary ray and include it into illumination model




Radiance for
refraction ray

$$I = I_E + K_A I_A + \sum_L (K_D (N \cdot L) + K_S (V \cdot R)^n) S_L I_L + K_S I_R + K_T I_T$$

Transparency

- Transparency coefficient is fraction transmitted
 - $K_T = 1$ if object is translucent, $K_T = 0$ if object is opaque
 - $0 < K_T < 1$ if object is semi-translucent




Transparency
Coefficient

$$I = I_E + K_A I_A + \sum_L (K_D (N \cdot L) + K_S (V \cdot R)^n) S_L I_L + K_S I_R + K_T I_T$$

Refractive Transparency

- For thin surfaces, can ignore change in direction
 - Assume light travels straight through surface




$$T \cong -L$$

Refractive Transparency

For solid objects, apply Snell's law:


$$\eta_r \sin \Theta_r = \eta_i \sin \Theta_i$$




$$T = \left(\frac{\eta_i}{\eta_r} \cos \Theta_i - \cos \Theta_r \right) N - \frac{\eta_i}{\eta_r} L$$

Recursive Ray Tracing

- Ray tree represents illumination computation



Ray traced through scene




Ray tree


$$I = I_E + K_A I_A + \sum_L (K_D (N \cdot L) + K_S (V \cdot R)^n) S_L I_L + K_S I_R + K_T I_T$$

Recursive Ray Tracing

- Ray tree represents illumination computation



Ray traced through scene



Ray tree

$$I = I_E + K_A I_A + \sum_L (K_D (N \cdot L) + K_S (V \cdot R)^n) S_L I_L + K_S I_R + K_T I_T$$

Recursive Ray Tracing

- **GetColor** calls **RayTrace** recursively

```
Image RayTrace(Camera camera, Scene scene, int width, int height)
{
    Image image = new Image(width, height);
    for (int i = 0; i < width; i++) {
        for (int j = 0; j < height; j++) {
            Ray ray = ConstructRayThroughPixel(camera, i, j);
            Intersection hit = FindIntersection(ray, scene);
            image[i][j] = GetColor(scene, ray, hit);
        }
    }
    return image;
}
```

Summary

- Ray casting (direct illumination)
 - Usually use simple analytic approximations for light source emission and surface reflectance
- Recursive ray tracing (global illumination)
 - Incorporate shadows, mirror reflections, and pure refractions

All of this is an approximation
so that it is practical to compute

More on global illumination later!

Illumination Terminology

- Radiant power [flux] (Φ)
 - Rate at which light energy is transmitted (in Watts).
- Radiant Intensity (I)
 - Power radiated onto a unit solid angle in direction (in Watts/sr)
 - e.g.: energy distribution of a light source (inverse square law)
- Radiance (L)
 - Radiant intensity per unit projected surface area (in Watts/m²sr)
 - e.g.: light carried by a single ray (no inverse square law)
- Irradiance (E)
 - Incident flux density on a locally planar area (in Watts/m²)
 - e.g.: light hitting a surface along a
- Radiosity (B)
 - Exitant flux density from a locally planar area (in Watts/ m²)